

# Transfer Between Saccade and Reach Adaptation Using Concurrent Sine-Wave Perturbations

جامعة نيويورك أبوظي

<sup>1</sup>New York University Abu Dhabi, <sup>2</sup>NYU Department of Psychology, <sup>3</sup>NYU Center for Neural Science, <sup>4</sup>NYUAD Center for Artificial Intelligence and Robotics, <sup>5</sup>NYUAD Center for Brain and Health

# NYU ABU DHABI

## Background


- Multiple studies suggest that saccade and reach planning share a common spatial representation:
- Reach accuracy improves with a saccade (even in darkness)<sup>1</sup>.
- Reach endpoints are consistently biased toward gaze location<sup>2</sup>.
- If both effectors use a common map, then systematic biases in one should also appear in the other.

### Question


If we adapt saccades and reaches concurrently, will we observe transfer of adaptation from one effector to the other?

### Procedure

- Both effectors adapted concurrently
- Perturbation magnitude follows a sinusoidal time course across trials<sup>3</sup>.




### Perturbation: Spatially orthogonal




## Perturbation: Temporally distinct


#### Predictions:



#### Results



In this example participant, transfer can be found from saccade to reach, but transfer from reach to saccade is in the opposite direction than predicted.



- Across participants, we found no significant amount of transfer between effectors.
- In individual conditions where transfer was significant, the direction of the transfer was not consistent across participants or conditions.

#### References

- 1. Bock, O. Exp Brain Res (1986).
- 2. Vazquez, Y. et al. *J Neurophysiol* (2017).
- 3. Hudson, T. E. and Landy, M. S. *J Neurosci Methods* (2012).

### Support

NIH EY08266 NYUAD Research Institute CG010, CG012

