

Contingent adaptation in masking and surround suppression



Hörmet Yiltiz¹, David J. Heeger^{1,2}, Michael S. Landy^{1,2} ¹Department of Psychology, ²Center for Neural Science, New York University

Introduction

Hebbian normalization model of adaptation¹: neurons that fire together more often inhibit each other more.

Hypothesis

Contingent adaptation: presenting stimuli synchronously leads to greater suppression compared to asynchronous presentation.

- Demonstrated in macague physiology².
- Untested in humans.

Conclusion

- Contingent adaptation raises detection threshold and changes appearance.
- Supports the Hebbian normalization model.

REFERENCE

- 1. Westrick et al., J Neurosci, 2016.
- 2. Aschner et al., SFN, 2016.


ACKNOWLEDGEMENTS Supported by NIH EY 08266.

CONTACT


hormet.viltiz@nvu.edu Available online: http://hyiltiz.com/VSS2018.pdf


DETECTION AND CROSS-ORIENTATION SUPPRESSION

- Adapt: non-contingent or contingent
- Measure: threshold contrast for detecting a masked grating



Which side has an extra grating?







Contingent adaptation results in stronger cross-orientation suppression and thus increased masked detection thresholds.

APPEARANCE AND SURROUND SUPPRESSION

- Adapt: vary center-surround contingency
- Measure: perceived matching contrast



Which center has higher contrast?



The center grating *appears* to have lower contrast when it matches the adapter, thus requires higher physical contrast to match the other center grating.